

ORIGINAL ARTICLE

INTUSSUSCEPTION IN CHILDREN: COMPARISON BETWEEN ULTRASOUND DIAGNOSIS AND OPERATION FINDINGS

Umesh Shah¹, Mukesh Kothari²

Author's Affiliations: ¹Assitant Professor, Dept. of Radiology; ²Associate Professor, Dept. of Surgery, GMERS Medical College, Dharpur, Patan, Gujarat

Correspondence: Dr Mukesh Kothari Email: drmukeshkothari@yahoo.com

ABSTRACT

Background: Intussusception is one of the more common causes of intestinal obstruction in children. The diagnosis may be based mainly on clinical features; however, there are no classic signs and symptoms that are common to all cases. This study reports our experience of diagnosis and operation findings of children with intussusceptions.

Materials and Methods: This was a retrospective review of intussusceptions in children in a tertiary health facility in a tropical developing country from January to December 2011.

Results: Twentyfive out of 41 children (M:F = 2.2:1) admitted with intussusceptions within the period were studied. The median age was 6.0 ± 5.57 months (range 3 months- 7 years). Ultrasonography positively diagnosed intussusceptions in 20 (80%) cases.

Conclusion: Ultrasonography can increase diagnostic confidence in intussusceptions.

Keywords: Children, intussusception, operation findings, tropical developing country, ultra-sonography

INTRODUCTION

Intussusception is one of the more common causes of intestinal obstruction in children. Recognising and treating this condition rapidly is important to prevent potentially fatal complications.^{1,2}

The diagnosis may be based mainly on clinical features,³ which could be quite challenging. This is, because, there are no classic signs and symptoms that are common to all cases of intussusception, a situation that often leads to delay in diagnosis. The classic triad of vomiting, abdominal pain and passage of blood per rectum occurs only in a third of cases.⁴⁻⁷

It implies that this classic triad cannot be wholly relied upon in making a diagnosis. Furthermore, atypical presentation of cases of intussusception is seen in up to 16% of children.^{8,9} Therefore, ultrasound scan (US) of the abdomen has been used to aid diagnosis and is said to be very reliable in experienced hands.^{10,11}

We sought to evaluate the value of Ultrasonography in increasing the diagnostic confidence in patients with suspected intussusceptions by deter-

mining the accuracy of Ultrasonography diagnosis and operation findings in cases of suspected intussusception in children in our centre. These findings may be useful to paediatric and other surgeons involved in the care of children with intussusception in similar settings as ours where late presentation is frequent and non-operative reduction is infrequently performed.

MATERIALS AND METHODS

The medical records of children aged 0-15 years consecutively admitted with intussusception to GMERS, Patan from January to December 2011 were retrospectively studied. The data extracted from the case notes included patient's age, sex, symptoms and signs on presentation in hospital, clinical, Ultrasonography and barium enema diagnoses, status of sonographer and operation findings. Data analysis was carried out with descriptive statistics using SPSS version 14 for windows. Children with clinical diagnosis of intussusception without Ultrasonography diagnosis as well as those who had prolapsed intussusceptions were excluded

from the study. Institutional consent was obtained from the Ethics committee. The sensitivity, specificity, positive and negative predictive values of the Ultrasonography scan were determined.

RESULTS

Forty-one patients with intussusception were seen during the 8years period. Ten of them did not have US diagnosis while six children had prolapsed intussusceptions and hence were excluded from the study. The age range at presentation was 3 months to 7 years with a median age of 6.0 ± 5.57 months. There were 21 (84%) infants and 4 (16%) older children made up of 17 (68%) males and 8 (32%) females (M:F = 2.13:1) [Table 1]. The peak incidence of intussusception was in the 3-7 months age group.

Table 2 summarises the sources of the Ultrasonography results. More than half ($n = 14$; 56%) of them were generated by private Ultrasonography outfits whose experiences in diagnostic Ultrasonography could not be ascertained. The remaining scans were carried out in the Radiology Department by resident medical officer in 8 (32%) and consultant radiologists in 3 (12%) patients respectively. The Ultrasonography reports produced by the resident doctors were also reviewed by their consultants who countersigned them. All the 11 (100%) reports generated from the Teaching Hospital became truly positive at the end, one, which also had double contrast barium enema, which diagnosed intestinal polyp as cause of the intussusception. Of the 14 (56%) results from the private outfits, 9 (64.3%) were truly positive for intussusception. None in this patient group had barium enema study. In all, there were 21 (84%) positive Ultrasonography results and 4 (16%) negative Ultrasonography results.

Out of the 21 positive results, 20 (80%) of them were truly positive for intussusception at operation while 1 (4%) was falsely positive [Table 3]. Here, an obstructed and dilated segment of bowel caused by an adhesive band produced Ultrasonography features of intussusception leading to misdiagnosis of the actual condition. Of the four negative results, 3 (12%) of them were falsely negative meaning that there was intussusception at operation while the remaining 1 (4%) was truly negative, meaning that there was no intussusception at operation. The total accuracy of Ultrasonography was 84% while the sensitivity and specificity were 87% and 50% respectively. The low specificity was probably due to inability to identify all of the patients who were referred and subsequently were proven not to have an intussusception. Similarly, the positive and negative predictive values were 95.2% and 25% respectively.

Table 1: Age and sex distribution of children with intussusception

| *Age | Sex | | Total (%) |
|-------------|----------|------------|-----------|
| | Male (%) | Female (%) | |
| 0-3 months | 3 (12) | 0 (0) | 3 (12) |
| 4-7 months | 8 (32) | 6 (24) | 14(56) |
| 8-11 months | 2 (8) | 2 (8) | 4 (16) |
| 1-2 years | 2 (8) | 0 (0) | 2 (8) |
| 3-7 years | 2 (8) | 0 (0) | 2 (8) |

* Age as at last birthday

Table 2: Distribution of the sources of US results

| Source | No. (%) |
|----------------------------|---------|
| Private Ultrasound Outfits | 14 (56) |
| Resident Doctors | 8 (32) |
| Consultant Radiologist | 3 (12) |

Table 3: Ultrasound diagnosis of intussusception compared with operation findings

| US Result | Intra operative diagnosis of intussusception | | | |
|-------------|--|------------|------------|----------------|
| | Present | | | Absent |
| | Ileo Caecal | Ileo Colic | Colo Colic | Adhesive Obstn |
| Positive(a) | 4 | 15 | 1 | (b) 1 |
| Negative(c) | 1 | 2 | 0 | (d) 1 |

Ileocolic ($n = 17$; 68%) and ileocaecal ($n = 5$; 20%) intussusceptions were more commonly encountered at operation. There was no entry of cases of intussusceptions that had spontaneously reduced in the case notes. The causes of the intussusceptions were unknown in 23 (92%) patients as no specific lead points were found at operation. How-

ever, in 2 (8%) other older patients a sessile polyp and an enlarged hypertrophied ileal lymphoid patch were documented as causes of the intussusceptions leading to resection of the bowel segments.

DISCUSSION

The results of this study showed that infants were those commonly diagnosed with intussusception. This finding was in agreement with those of other studies.¹²⁻¹⁵ It differed with earlier reports that associated childhood intussusception with the above 5 years age group.^{16,17} This underscores the challenge in diagnosis that may be experienced.

While the older child can appreciate his symptoms to some extent, the preverbal children cannot give any clue about their problem. Therefore, total reliance must be placed on the parents' history and the objective assessment of the surgeon. However, there are no classic signs and symptoms that are common to all cases of intussusception. Even the classic triad of vomiting, abdominal pain and passage of blood per rectum is said to occur only in a third of cases.⁴⁻⁷ Recent studies have it that about 30%-68% of children with clinical findings suggestive of intussusceptions end-up with the condition.^{18,19}

This underscores the need for an additional diagnostic tool such as Ultrasonography scan, which has been said to be very reliable in experienced hands.^{10,11} The finding of the study showed that majority of the patients had their Ultrasonography investigations performed by private Ultrasonography outfits. The importance of this fact is that, the eventual result will depend on the experience of the sonographer.^{10,11} This is likely to influence the accuracy of the investigation. Presumably, there may also be patients referred to the centre who had not been identified to be included in the study.

On the other hand, the scans performed by the Resident doctors were again reviewed by the consultants who were more experienced. This led to a 100% positive predictive value of the children with intussusception. On the contrary, higher frequency Ultrasonography imaging with better resolution were more likely to be available for use in specialist Teaching Hospital setting¹⁹ than on private outfits.

The results of the study showed that Ultrasonography was highly accurate in the diagnosis of intussusception in experienced hands with a sensitivity of 87%. This is in line with earlier series in the developed economies.¹⁹⁻²¹ However, the observed specificity of 50% appeared low and at variance with 88%-100% normally quoted in literatures.^{10,18,19} Presumably this low value may be due to patients referred to our centre who were not identified to be included in the study. It could also be accounted for by the inexperience of the private sonographers who diagnosed a child with adhesive

bands as having intussusception, leading to one false positive result. This agrees with the observation that thickened bowel wall in some acute gastrointestinal conditions in children may produce Ultrasonography features of intussusceptions.²² This therefore calls for experience in Ultrasonography interpretation and a high index of suspicion on the part of the clinician. Where still in doubt, radiography with barium continued into non-operative management of the condition is performed except in late presentation with advanced symptoms, such as peritonitis indicating the presence of gangrenous intestine.

To perform barium enema reduction, the patient must be adequately prepared and theatre booked and ready for operative treatment should the procedure fail. A column of barium is retained at a level of 30 cm above the buttocks of the patient.^{23,24} A higher level may result in an increased intraluminal pressure with risk of perforation due to impaired viability of the intestine. Reduction is considered complete when barium refluxes freely into the small intestine for more than 5-10 cm. If this fails or if perforation occurs, immediate surgery is performed.

The realized accuracy rate of 84% was much lower than 94.4% reported by other series.^{19,22} While relatively high positive predictive value comparable to those in published series was achieved, the negative predictive value was rather low and not in keeping with those of other studies.^{19,22} The types of intussusceptions found at operation also agreed with those commonly reported in children in our environment and world-wide.⁴⁻⁷ Similarly, the causes of intussusceptions documented in the study were in keeping with those of other series.^{25,26}

In conclusion, the accuracy of Ultrasonography diagnosis in experienced hands was high and could increase the diagnostic confidence in patients with suspected intussusceptions. Thickened bowel wall in other gastrointestinal conditions could produce Ultrasonography features of intussusceptions. Therefore, a high index of suspicion is recommended. If still in doubt after Ultrasonography diagnosis, we recommend a contrast enema, which can be continued into non-operative management. However, in the tropical developing countries, operative treatment may sometimes be expedient in averting grave consequences of the condition, which could be occasioned by poverty or non-availability of the required non-operative modality of treatment.

REFERENCES

1. Fallat ME. Intussusception. In: Ashcraft KW, editor. Pediatric Surgery. 3rd ed. W.B. Saunders Company Philadelphia, Pennsylvania; 2000. p. 518-26.
2. Young DG. Intussusception. in: O'Neil JA Jr, Rowe MI, Grosfeld JL, Fonkalsrud EW, Coran AG, editors. Pediatric Surgery. 5th ed. Mosby, New York, USA. 1998. p. 1185-98.
3. Dennison WM. Acute intussusception in infancy and childhood. Glasgow Med J 1948;29:71-80.
4. Stringer MD, Pablot SM, Brereton RJ. Paediatric intussusception. Br J Surg 1992;79:867-76.
5. Bruce J, Borzi PA. Intussusception in childhood. Surgery 1992;10:213-6.
6. Mangete ED, Allison AB. Intussusception in infancy and childhood: An analysis of 69 cases. West Afr J Med 1994;13:87-90.
7. Ugwu BT, Legbo JN, Dakum NK, Yiltok SJ, Mbah N, Uba FA. Childhood intussusception: A 9-year review. Ann Trop Paediatr 2000;20:131-5.
8. Baracchini A, Chiaravallotti G, Quinti S, Rossi A, Favili T, Ughi C, *et al.* Intestinal intussusception in children. Minerva Pediatr 1995;47:215-9.
9. Shekhawat NS, Prabhakar G, Sinha DD, Goyal RB, Gupta A, Sharma RK, *et al.* Nonischemic intussusception in childhood. J Pediatr Surg 1992;27:1433-5.
10. Shanbhogue RL, Hussain SM, Meradji M, Robben SG, Vernooij JE, Molenaar JC. Ultrasonography is accurate enough for the diagnosis of intussusception. J Pediatr Surg 1994;29:324-8.
11. John SD. The value of ultrasound in children with suspected intussusception. Emerg Radiol 1998;5:297-305.
12. Adejuyigbe O, Jeje EA, Owa JA. Childhood intussusception in Ile-Ife, Nigeria. Ann Trop Paediatr 1991;11:123-7.
13. Archibong AE, Usoro IN, Ikpi E, Inyang A. Paediatric intussusception in Calabar, Nigeria. East Afr Med J 2001;78:19-21.
14. Carneiro PM, Kisusi DM. Intussusception in children seen at Muhimbili National Hospital, Dar es Salaam. East Afr Med J 2004;81:439-42.
15. Boudville IC, Phua KB, Quak SH, Lee BW, Han HH, Verstraeten T, *et al.* The epidemiology of paediatric intussusception in Singapore: 1997 to 2004. Ann Acad Med Singapore 2006;35:674-9.
16. Joly BM, Thomas HO. Non-infantile idiopathic intussusception in Western Nigeria; with a report of 33 cases. West Afr Med J 1954;3:3-16.
17. Elebute EA, Adesola AO. Intussusception in western Nigeria. Br J Surg 1964;51:440-4.
18. Pracros JP, Tran-Minh VA, Morin de Finfe CH, Defrenne-Pracros P, Louis D, Basset T. Acute intestinal intussusception in children. Contribution of ultrasonography (145 cases). Ann Radiol (Paris) 1987;30:525-30.
19. Verschelden P, Filiatrault D, Garel L, Grignon A, Perreault G, Boisvert J, *et al.* Intussusception in children: Reliability of US in diagnosis – A prospective study. Radiology 1992;184:741-4.
20. Bhisitkul DM, Listernick R, Shkolnik A, Donaldson JS, Henricks BD, Feinstein KA, *et al.* Clinical application of ultrasonography in the diagnosis of intussusception. J Pediatr 1992;121:182-6.
21. Eshed I, Gorenstein A, Serour F, Witzling M. Intussusception in children: Can we rely on screening sonography performed by junior residents? Pediatr Radiol 2004;34:134-7.
22. Cina M, Rahim F, Davudi M. The accuracy of ultrasonography technique in detection of the intussusception. J Appl Sci 2009;9:3922-6.
23. Girdany BR, Bass LW, Sieber WK. Roentgenologic aspects of hydrostatic reduction of ileocolic intussusception. Am J Roentgenol Radium Ther Nucl Med 1959;82:455-61.
24. Saxton V, Katz M, Phelan E, Beasley SW. Intussusception: A repeat delayed gas enema increases the nonoperative reduction rate. J Pediatr Surg 1994;29:588-9.
25. Kong FT, Liu WY, Tang YM, Zhong L, Wang XJ, Yang G, *et al.* Intussusception in infants younger than 3 months: A single center's experience. World J Pediatr 2010;6:55-9.
26. Uddin W, Kella N, Ahmed J, Mehmood T. An audit of non- idiopathic intussusceptions in children. J Liaquat Univ Med Health Sci 2010;9:134-7.