

ORIGINAL ARTICLE

BREAST LUMPS IN A TEACHING HOSPITAL: A 5 YEAR STUDY

Chiragkumar L Prajapati¹, Rajesh kumar K Jegoda¹, Upendra A Patel², Jay Patel³**Authors' Affiliation:** ¹Assistant professor; ²Associate professor, General Surgery, GMERS Medical College, Patan, Gujarat; ³MBBS, Patan, Gujarat**Correspondence:** Dr. Chiragkumar L Prajapati, Email: drchirag80@gmail.com

ABSTRACT

Background: Breast lump is the most common reason for presenting to surgery department. It is a source of great anxiety to a female when it is discovered. This study aims at a clinico-pathologic review of breast lump as a presenting complaint.**Material and Methods:** A 5-year retrospective analysis of 550 patients presenting with a complaint of breast lump to the surgery department of a Teaching Hospital in Western India from January 2008 December 2012.**Results:** Breast lump was the presenting complaint in 550(82.8%) of 664 patients, the lump was painless in 458 (83.3%), associated with pain and breast ulcer in 59(10.7%) and 33(6.0%) patients respectively; no palpable lump in 23 patients (4.2%) and only 54 patients (9.7%) had >1 lump.

The clinical diagnosis were breast cancer in 260 patients (47.3%), fibroadenoma in 175 (31.8%), fibrocystic changes in 67 (12.2%) patients; the others were benign diseases. Histopathology, done in 294 patients, revealed 161(54.8%), 56(19.0%) and 46(15.6%) patients having invasive cancer, fibroadenoma, and fibrocystic changes respectively.

Conclusion: Breast lump was the most common presenting complaint with most patients not presenting early. Fewer lumps are discovered by breast self examination. The finding of fibroadenoma as the most common of the benign lesions is similar to that reported by other researchers in India and other parts of the world.**Keywords:** breast lump, fibroadenoma, benign, malignant

INTRODUCTION

Breast lump is the most common reason for presenting to surgery departments. Most of these patients, however, are in a state of heightened anxiety until they have undergone specialist assessment, the necessary investigations and eventual reassurance. The majority of patients referred to a surgery department is said to have benign disease.¹⁻³ Fibroadenoma is the most common of the benign breast diseases.^{4,7} This study aims to audit the clinicopathologic features of patients with breast lump at teaching hospital in western India.

histology diagnosis for patients who returned with histology report were extracted from the case files.

Statistical Package for Social Sciences (SPSS) version 15.0 was used for data analysis. Simple frequencies were determined for variables.

RESULTS

Breast lump was the presenting complaint in 550(82.8%) of 664 patients presenting to the surgery department.

METHODOLOGY

The data on all patients who presented with history of breast lump to the surgical outpatient clinic at a teaching hospital in western India from 1st January 2008 to 31st December 2012 were retrospectively analysed. Information on age at presentation, parity, duration of symptom before presentation, mode of discovery of lump, previous breast disease, side and quadrant of breast affected, maximum diameter of breast lump, clinical diagnosis made by the senior registrar or unit consultant, refusal or acceptance of biopsy, type of biopsy done and

Table 1: Age wise distribution of subjects

Age in Years	Frequency
<20	59
21 – 30	148
31 – 40	130
41 – 50	115
51 – 60	60
61 – 70	22
71 – 80	16
Total	550

Females constitute 548(99.6%) with only 2(0.4%) males. The mean age of patients presenting with breast lump was 37.5 years [Range 13-80years]. Five hundred and forty-six (99.3%) were Igbos, as shown in table 1.

Table 2: Clinical Diagnosis of subjects

Clinical Diagnosis	Frequency (%)
Breast Cancer	260 (47.3)
Fibroadenoma	175 (31.8)
Fibrocystic disease	67 (12.2)
Normal Breast	8 (1.5)
Breast abscess	6 (1.1)
Duct ectasia	5 (0.9)
Galactocele	5 (0.9)
Inflammatory mass	4 (0.7)
Mastitis	4 (0.7)
Phyllodes Tumor	3 (0.5)
Mondor's disease	2 (0.4)
Recurrent Cancer	2 (0.4)
Kaposi Sarcoma	1 (0.2)
Others	8 (1.5)
Total	550 (100.0)

423 (77.2%) were premenopausal and 2 (0.4%) premenstrual. Nulliparity was seen in 208 (44.2%). The lump was painless in 458 (83.3%), associated with pain and breast ulcer in 59(10.7%) and 33(6.0%) patients respectively. Duration from discovery of lump to presentation was < 1 months in 102 (18.5%), 1-3 months 98 (17.8%), 3-6 months 73 (13.3%), 6-12 months 71 (12.9%), > 12 months in 192 (35%) and ranges from 2 days to 19 years. The duration was not documented in 14 patients.

The lumps were discovered by breast self examination (BSE) in 79 (14.3%) patients, they were accidental discovery in 293 (53.3%) patients. Attention was drawn to the lump by breast pain in 66 (12.0%), while the discovery was made by the doctor and the patients husband in 21 (3.8%) and 2 (0.4%) respectively. Mode of discovery was not stated in 89 (16.2%) patients. The lumps were increasing in size in 223 (49.6%), decreasing in 17 (3.8%), stationary in 178 (38.4%), fluctuating in 9 (2%).

The left breast was slightly more affected 262 (48.2%) than the right 252 (46.3%). The upper outer quadrant is the most common site of occurrence 165 (30%) followed by the upper inner quadrant 90 (16.4%), lower outer quadrant 50 (9.1%), central 44 (8%), whole breast 41 (7.5%), multiple quadrant 33 (6%), lower inner quadrant 19 (3.5%), no palpable lump 23 (4.2%) and in 85 (15.5%) patients the involved quadrant was not stated. Multiple lumps were found in 54 (9.7%) patients. The mean diameter of the lumps was 6cm [Range 0.5-30cm]. The number of patients admitting to previous history of breast disease was 43, with 30 having benign breast lesion. The clinical diagnosis (Table 2)

Were breast cancer in 260 patients (47.3%), fibroadenoma in 175 (31.8%), fibrocystic changes in 67 (12.2%) patients. The others were different types of benign diseases.

Histopathology Diagnosis

Biopsy was done on 346(62.9%) patients, 165(30%) refused biopsy, and 39(7.1%) patients did not require biopsy. The types of biopsies done were excision in 181 (52.3%) patients, 155 (44.8%) had incision biopsy, 5 (1.45%) core-cut biopsy, and 5(1.45%) FNAC. Histopathology report (Table 3) was available in 294 patients, with 161(54.8%), 56(19.0%) and 46(15.6%) patients diagnosed respectively as having invasive cancer, fibroadenoma, and fibrocystic changes. However, on clinical ground benign breast diseases constitute 53.3% (293 cases) with malignant being 46.7%. The mean age for breast cancer was 46.3 years, median 45 years range 26-78 years with a peak of 50 years. For fibroadenoma, mean age was 23.2 years, median 22years, range 18-39 years with a peak age of 21 years and for fibrocystic changes, mean was 37, median 35.5, range 16-80 years with peak age of 32 years. The breast lumps ranges in size from 0.5-30 cm, 0.5-20 cm, 0.5 - 11cm for breast cancer, fibroadenoma and fibrocystic changes respectively.

Table 3: Histopathological Diagnosis

Clinical Diagnosis	Frequency (%)
Breast Cancer	161 (54.8)
Fibroadenoma	56 (19.0)
Fibrocystic disease	46 (15.6)
DCIS	3 (1.0)
Fat necrosis	4 (1.4)
Phyllodes (benign)	4 (1.4)
Phyllodes (borderline)	1 (0.3)
Phyllodes (malignant)	1 (0.3)
Tuberculous Mastitis	3 (1.0)
Adenosis	3 (1.0)
Abscess	3 (1.0)
Tubular adenoma	2 (0.7)
Others	7 (2.4)
Total	294 (100.0)

DISCUSSION

The fear that a breast lump might be cancerous makes patient to present to the hospital. Fortunately, studies have shown that majority of patients presenting to a surgery department have benign disease. Breast lump was the most common presenting complaint (82.8%) to the symptomatic surgery department.¹⁻³ Most of these patients still present late to the hospital with 81.5% presenting after a month of noticing the lump.

Okobia et al⁸ noted that 78% of their patients reported after 3 months of symptoms while Atoyebi et al⁹ noted that 64% of their patients presented after 6 months, earliest been at 2weeks in a nurse.

Only small proportions (14.3%) of patients discover their lump by BSE, in majority (53.3%) breast lump was an accidental discovery. Jebbin et. Al¹⁰. working in Port-Harcourt noted that 20% of their study population performs BSE monthly. Following breast biopsies, histopathology revealed that breast cancer (54.8%) is slightly

commoner than benign breast disease (45.2%). The percentage of benign cases in our series is lower than 72.6%, 90%, 71.3% and 73% reported respectively in Eku¹¹, Benin¹², Zaria¹³ and Kano¹⁴ recently. But it compares with the 48% reported in Ibadan by Ajayi et al¹⁵ more than 30 years ago. When Ibadan was one of the very few centres handling such cases. The higher rate of malignant breast lesion in this series may be due to proliferation of private hospitals in our environment with the implication that benign breast lesion are treated there while malignant cases are referred to specialized surgery department like ours.

The most common cause of breast lumps in this series remained breast cancer (54.8%), followed by fibroadenoma (19.0%) and fibrocystic changes (15.6%). Other researchers have noted fibroadenoma to be the most common cause of breast lump^{2,13,16,17}, followed by fibrocystic disease, then breast cancer² or breast cancer then fibrocystic disease⁴. In London Ellis¹⁸ from a referral hospital noted that breast cancer was the most common finding in breast biopsies just like in our center. The age range of patients with breast lump in our study was 13-80 years. Eltahir et al¹⁹ found the age range to be 15-91 year in their patients with breast lumps. Thus it appears that patients with breast lumps in our environment are slightly younger.

The mean age for breast cancer in our environment had been noted to be 46.2 years²⁰, our study found 46.3 years which is very consistent with this previous finding. In the study by Carty et al²¹ on fibroadenoma, the age range of their study population was 15-48 years with a mean age of 28-years. But our study noted 18-39 years age range with a mean of 23.2years, this is consistent with other reports^{11,12,14,22}.

Of the minor breast lesions, this study found 6 (2%) cases of phyllodes tumour and 3 cases of tuberculous mastitis. The series by Anyanwu²³ 20 years ago reported only a case of phyllodes tumour but no case of tuberculous mastitis was reported. Also Akhator¹¹ reported only one case of phyllodes tumour with two cases of tuberculous mastitis. Though our study reported greater number of phyllodes tumour, this condition is still rare and is still within the 2%-4.4% rate reported.²⁴

CONCLUSION

Our study shows that breast lump was the most common presenting complaint with most patients not presenting early. Fewer lumps are discovered by breast self examination. The finding that breast cancer was slightly more common than benign breast lesions is at variance with most studies probably due to proliferation in our environment of private owned hospitals by general practitioners where most of the benign lesions are probably managed. But the finding of fibroadenoma as the most common of the benign lesions is similar to that reported by other researchers in India and other parts of the world.

REFERENCES

- Osime OC, Ohanaka EC. Analysis of five-year breast biopsies carried out in the University of Benin Teaching Hospital Benin City. *Niger PostgradMed J.* 2008 ;15(3):160-163.
- Ohene-Yeboah MO. An audit of excised breast lumps in Ghanaian women. *West Afr J Med.* 2005;24(3):252-255.
- Clegg-Lamptey JN, Aduful HK, Yarney J, Adu-Aryee NA, Vanderpuye V, Kyereh M, et al. Profile of breast disease at a self-referral clinic in Ghana. *NepalMedColl J.* 2004;6(2):129-132.
- Oluwole SF, Freeman HP. Analysis of benign breast lesions in blacks. *Am J Surg.* 1979 Jun;137(6):786-789.
- Sönmez K, Türkyılmaz Z, Karabulut R, Demirođullari B, Ozen IO, Moraliödlü S, Surgical breast lesions in adolescent patients and a review of the literature. *Acta Chir Belg.* 2006 Jul-Aug;106(4):400-404.
- Adeniji KA, Adelusola KA, Odesanmi WO. Benign disease of the breast in Ile-Ife: a 10 year experience and literature review. *Cent Afr J Med.* 1997;43(5):140-143.
- Ihekwa FN. Benign breast disease in Nigerian women: a study of 657 patients. *J R Coll Surg Edinb.* 1994;39(5):280-283.
- Okobia MN, Ahigbe JU. Pattern of malignant disease at University of Benin Teaching Hospital. *TropDoct* 2005;35(2):9-12.
- Atoyebi OA, Atimomo CE, Adesanya AA, Beredugo BK, da Rocha-Afodu JT. An appraisal of 100 patients with breast cancer seen at the Lagos University Teaching Hospital. *Nig. Qt. J Hosp.Med.* 1997;7(2):104-108.
- Jebbin NJ, Adotey JM. Attitudes to, knowledge and practice of breast self-examination (BSE) in PortHarcourt. *Nig J Med* 2004;13(2):166-170
- Akhator A. Benign breast masses in Nigeria. *Nig J Surg.* 2007;17(2):105-108
- Okobia MN, Osime U. A clinicopathological study of benign breast disease in Benin City, Nigeria. *Nig J Surg.* 1998;5(2):64-69
- Yusuf LMD, Odigie VI, Mohammed A. Breast masses in Zaria, Nigeria. *Annals of African Medicine.* 2003;2(1):13-16.
- Ochicha O, Edino ST, Mohammed AZ, Amin SN. Benign breast lesions in Kano. *Nig J of SurgicalResearch.* 2002;4:1-5.
- Ajayi OO, Adekunle O. Non-malignant breast masses in an African population. *Br. J. Surg.* 1973;60:465-468
- Adesunkanmi AR, Agbakwuru EA. Benign breast disease at Wesley Guild Hospital, Ilesha, Nigeria. *West Afr J Med.* 2001;20(2):146-151
- Otu AA. Benign breast tumours in an African population. *JRColl Surg.* 1990;35(6):373-375
- Ellis H. Review of general surgery 1984-85. *Postgraduate Medical Journal.* 1985; 61: 941- 956
- Eltahir A, Jibril JA, Squair J, Heys SD, Ah-See AK, Need-Ham G, et al. The accuracy of "onestop" diagnosis for 1 110 patients presenting to a symptomatic breast clinic. *J R Coll Surg Edinb.* 1999; 44:226-230
- Anyanwu SNC. Temporal trends in breast cancer presentation in the third world. *J Exp Clin CancerRes.* 2008; 27(1): 17
- Carty NJ, Carter C, Rubin C, Ravichandran D, Royle GT, Taylor I. Management of fibroadenoma of the breast. *Ann R Coll Surg Engl* 1995; 77: 127-130
- Anyanwu SNC. Fibro-adenoma of the breast in Nigerian Igbo. *SAfrMed J.* 2000;90(12):1223- 1226
- Anyanwu SNC. Spectrum of breast disease among Nigerian Igbo. *Orient Journal of Medicine.* 1990;2(1):14-15
- Petrek JA. Phyllodes tumours. In Harris JR, Lippman ME, Morrow M, Osborne CK, ed. *Harris disease of breast. 2 Ed.* Lippman, LippincottWilliams andWilkins;2000:47.