

ORIGINAL ARTICLE

THE SUTURELESS CIRCUMCISION- AN ALTERNATIVE TO THE STANDARD TECHNIQUE

Hiren D Parmar¹, Supreet D Bhatt²¹Assistant Professor; ²Resident, Department of Surgery, Smt NHL Municipal Medical College, Ahmedabad

Correspondence: Dr. Hiren D Parmar, Email: drhirenparmar@gmail.com

ABSTRACT

Introduction: circumcision is a commonly performed surgery. Surgeons have become increasingly interested in the use of adhesive bonds. Recent advances have been made in the use of tissue glue in the circumcision. In this study, we used 2-octyl cyanoacrylate for closing circumcision wounds and we have reported our experience.

Material & methods: this was a prospective non-comparative preliminary clinical study involving 30 patients where 2-octyl cyanoacrylate was used as a tissue adhesive for wound closure after formal circumcision at surgery department from May 2008 to Nov 2010.

Result: in this study, the mean time taken for skin closure by 2-octyl cyanoacrylate is much faster in last year of study. There is significant less pain. There is 10% (3 cases) of complications are observed. The wound cosmesis score on 90th post-operative day is optimal except in one patient of wound separation.

Conclusion: the comparison with criterions of time taken for skin closure, the post-operative pain, the cosmetic appearance of adhesive glue proves that 2-octyl cyanoacrylate skin closure is significantly better than the traditional skin suturing skin closure.

Keywords: 2-octyl cyanoacrylate, suture less circumcision, adhesive glue skin closure

INTRODUCTION

The circumcision is one of the most commonly performed operations, with various recognised techniques. Male circumcision is the removal of some or the entire foreskin (prepuce) from the penis¹. Procedures include the plastibell with ligature², gomco clamp³, mogen clamp⁴ and excision with or without suturing. In last 20 years, surgeons have become increasingly interested in the use and application of adhesive bonds. Recently advances have been made in the use of tissue glue in circumcision⁵. By undertaking this study, we begun using 2- octyl cyanoacrylate⁶ for closing circumcision wounds and here in we have reported our experience. Our aims & objectives are to study if circumcision wound closure is feasible with topical skin adhesive (2- octyl cyanoacrylate), to study complications, cost effectiveness, merits & demerits of the procedure and to assess if the technique can be used as reasonably good alternative to conventional suture circumcision.

MATERIAL & METHODS

This was a prospective non-comparative preliminary clinical study involving 30 patients where 2- octyl cyanoacrylate was used as a tissue adhesive for wound

closure after formal circumcision. 30 healthy male patients from the age of 1 year to 75 years, during a period of May 2008 to Nov 2010 in our surgical department, Smt S.C.L. Municipal Medical College comprised the study group. There were no exclusion criteria. The indications for surgery⁷ were chronic balanoposthitis, congenital or acquired phimosis, paraphimosis, long prepuce skin, religious reason and others like zip injury. Appropriate blood tests were performed and consent for surgery was taken. The operation was performed under local anaesthesia (dorsal penile block⁸) in the elderly patient while general anaesthesia in the younger patient. The technique of dorsal slit followed by free hand cutting all around with sharp scissors was used in all the cases⁹. The outer layer of the foreskin was retracted back and meticulous haemostasis was achieved. Incision was cleaned after it. The cut edges were approximated with forceps and the glue 2-octyl cyanoacrylate was applied in two thin layers¹⁰. In some cases suture with 3-0 chromic catgut placed 90 degree apart helped the application of the glue. The glue was allowed to harden and polymerized. Leakage of the glue between the edges was avoided so that hardened glue does not catch the undergarments. After the procedure was finished, the wound was dried and the time of start of skin closure and the time of finishing the skin closure were

noted down using a stopwatch timer. The time taken for skin closure was noted. No liquid or antibiotic ointment applied after glue application. Protective dry gauze applied after adhesive film was completely solid/polymerized (approximately 5 minutes after application). All the patients received a 5 day course of ampicillin+cloxacillin and analgesic ibuprofen & paracetamol in appropriate doses. Adult were also given oral oestrogen tablets.

All the patients were discharged on the same day and followed on the 1st, 5th, 10th, 15th, 30th and 90th day. Bathing on the operative site was permitted after the 5th day onwards. Post-operative pain was assessed at 1st day, 5th day, 10th day, 15th day, 30th day and 90th day using Visual Analogue Score¹¹ of 0 to 100, 0 being no pain and 100 being worst pain possible assessed by patients themselves. All wounds were assessed by visual inspection at 1, 5, 10 days after wound closure. Wound was scored by using wound ASEPSIS score¹².

Confirmatory culture was not routinely performed. The Modified Hollander Cosmesis Scale (mHCS)¹³, a validated scale, was used to evaluate at 15th, 30th, 90th day: (1) step-off borders (2) edge eversion (3) contour irregularities (4) excess inflammation (5) wound margin separation (6) overall appearance. A total cosmetic score was derived by adding the scores of variables. A score of 1 was given to each variable if present in the wound, so a score of 6 was considered as worst while 0 as best. Any complications or infections, if present were also observed.

RESULTS

The time taken for skin closure is measured using a stopwatch timer and entered in units of seconds. The skin closure time using 2- octyl cyanoacrylate is longer in first year and shorter in last year reflecting a learning curve associated with application.

Table 1: Time taken for skin closure

Time period	Patients	Minimum time (seconds)	Maximum time (seconds)	Average time (seconds)
May 2008 to April 2009	14	80	110	90
May 2009 to April 2010	7	60	80	70
May 2010 to Nov 2010	9	40	80	60

Table 2: Post-operative pain score

Time (day)	Average visual analogue score
1 st	50
5 th	10
10 th	0
15 th	0
30 th	0
90 th	0

A Visual Analogue Scale (VAS) is a measurement instrument that tries to measure a characteristic or

attitude that is believed to range across a continuum of values and cannot easily be directly measured. Operationally a VAS is usually a horizontal line, 100 mm in length, anchored by word descriptors at each end. The patient marks on the line the point that they feel represents their perception of their current state of pain. The VAS score is determined by measuring in millimetres from the left hand end of the line to the point that the patient marks. The score was never on higher side from 1st to 90th post-operative day.

Table 3: Wound asepsis score

Interval (days)	No complication	Seroma	Erythema	Purulent discharge	Wound separation	total
1	28 patients	2 patients	0	0	0	30
5	28 patients	1 patient	0	1 patient	0	30
10	29 patients	0	0	0	1patient	30

The points were taken in account: Additional treatment, the presence of Serous discharge, Erythema, Purulent exudate, and Separation of the deep tissues, the Isolation of bacteria, and the duration of inpatient

Stay (ASEPSIS). Overall 3 patients developed wound complications but wound asepsis score was not on higher side.

Table 4: Wound Cosmesis score

Time (day)	Score 0	Score 1	Score 2	Score 3	Score 4	Score 5	Score 6
15	27 patient	Nil	Nil	1 patient	1 patient	1 patient	Nil
30	28 patient	Nil	Nil	Nil	1 patient	1 patient	Nil
90	29 patient	Nil	Nil	Nil	Nil	1 patient	Nil

It was measured by using the modified Hollander Scale. The modified Hollander Scale assigns a score ranging

from 0 (best) to 6 (worst) for six categories. In most of the patients, it was 0. It indicates good cosmesis.

DISCUSSION

Now a day surgeons are looking for faster, comfortable and cosmetically best technique for skin closure, more over 2-octyl cyanoacrylate is easier to use and provides a flexible, water resistant, sealed skin closure¹⁴. 2-Octyl cyanoacrylate provides a needle-free method of wound closure, an important consideration because of blood-borne viruses (e.g. HIV)¹⁵. It requires no bandaging due to its antimicrobial properties¹⁶. For the patient side, it gives less pain during post-operative period, needs no suture or staple removal, disappears naturally as incision heals, leaves no marks and patients can even have a shower¹⁷.

In the present study, most of the patients (53.33%) were in age group of 1-10 years. It was not significant as patients were selected randomly but it can be concluded that circumcision is generally performed in age group 1-10 years. Circumcision was most commonly done for congenital phimosis followed by acquired phimosis (66.67%)¹⁸. In one of the first published studies evaluating octylcyanoacrylate, Quin.J.et.al.¹⁹ indicates that use of the skin adhesive is found to be significantly faster. In Matin S.F. study²⁰ & in study by James M. Elmore, Edwin A. Smith, and Andrew J.Kirsch²¹ it is to be concluded that the skin adhesive technique is significantly faster. In present study, the mean time taken for skin closure in adhesive glue is much faster in last year of the study. This is of great significance as it is suggestive that glue application requires practice and as experience is gained application becomes easier and faster. In the earlier studies Zempsky W.T. et.al.²² and Quinn J. et.al.¹⁹ have compared the post-operative pain visual analogue scale of 0 to 100 and have shown less post-operative pain in adhesive glue group. In the present study, there is less post-operative pain with adhesive glue up to first 5 days following surgery from 10 days onwards there is no pain. Singer A.J. et.al.²³ shows that at the end of 1st week after surgery were similar and fewer cases of adhesive glue were erythematous. Wound dehiscence rate is 1.6% in adhesive glue group and 0.9% in suturing group. In Toriumi D.M. et.al.²⁴ they had evaluated wound at 1st week and had not observed any complication. In study Suture less Circumcision Using 2-octyl cyanoacrylate (Dermabond): appraisal after 18 months of experience by James M. Elmore, Edwin A. Smith, and Andrew J. Kirsch²¹ no patient developed wound complication. In present study, 10% (3 cases) of complications are observed. There are two seroma (1 purulent) and 1 wound separation observed. The study conducted by Toriumi D.M.²⁴ et.al. Observed wounds using Modified Hollander Cosmesis Scale and later by Visual Analogue Scale revealed the equivalent result with formal scale and superior result with later scale for wound cosmesis. In study done by Jallali N. et.al.²⁵ showed no significant difference in cosmesis with both the score. In present study, wound cosmesis score on post-operative 90th day is optimal except in 1 patient of wound separation.

In comparison with criterions of time taken for skin closure, the post-operative pain, the cosmetic appearance of adhesive glue, 2-octyl cyanoacrylate skin closure is significant better than the traditional skin suturing skin closure²¹.

REFERENCES

1. Alanis MC, Lucidi RS. Neonatal circumcision: a review of the world's oldest and most controversial operation. *Obstet Gynecol Surv.* 2004;59:379-95.
2. Khairi A., El-Kholi N., Tolba M., Ismail T., Dwaba M., Hafez M., *et al.* plastibell circumcision; evaluation of a technique in 800 cases. *Egyptian Journal of Surgery* 2005;1:1
3. David Peleg, Ann Steiner. The Gomco Circumcision: Common Problems and Solutions. *Am Fam Physician* 1998;58:891-8.
4. Reynolds RD. Use of the Mogen clamp for neonatal circumcision. *Amer Fam Physician* 1996;54:177-82.
5. Cheng W, Saing H. A prospective randomized study of wound approximation with tissue glue in circumcision in children. *J Paediatr Child Health* 1997;33:515-6.
6. Bruns TB, Worthington JM. Using tissue adhesive for wound repair: a practical guide to dermabond. *Am Fam Physician* 2000;61:1383-8.
7. A.M.K. Rickwood. Medical indications for circumcision. *BJU International* 1999;1:45-51
8. Telgarsky B, Karovic D, Wassermann O, Ogbovicova E, Csomor D, Koppl J, *et al.* Penile block in children, our first experience. *Bratisl Lek Listy* 2006;107:320-2.
9. Holman JR, Lewis EL, Ringler RL. "Neonatal circumcision techniques". *American Family Physician* 1995;52:511-8, 519-20.
10. Kelly BD, Landon DJ, Timlin ME, Sheikh M, Nusrat NB, D'Arcy FT, *et al.* Paediatric sutureless circumcision--an alternative to the standard technique. *Pediatr Surg Int* 2012;28:305-8.
11. Bodian, Carol A., Freedman, Gordon, Hossain, Sabera, *et al.* The Visual Analogue Scale for Pain: Clinical Significance in Postoperative Patients. *Anesthesiology* 2001:1356-61
12. Wilson AP, Weavill C, Burrige J, Kelsey MC. The use of the wound scoring method 'ASEPSIS' in postoperative wound surveillance. *J Hosp Infect* 1990;16:297-309
13. Zhang Zhong-tao, Zhang Hong-wei, Fang Xue-dong, Wang Li-ming, Li Xiao-xi, Li Ya-fen, *et al.* Original article Cosmetic outcome and surgical site infection rates of antibacterial absorbable (Polyglactin 910) suture compared to Chinese silk suture in breast cancer surgery: a randomized pilot research. *Chinese Medical Journal* 2011;124:719-24
14. Singer AJ, Thode HC Jr. A review of the literature on octylcyanoacrylate tissue adhesive. *Am J Surg* 2004;187:238-48.
15. Kaye JD, Kalisvaart JF, Cuda SP, Elmore JM, Cerwinka WH, Kirsch AJ. Suture less and scalpel-free circumcision--more rapid, less expensive and better? *J Urol* 2010;184:1758-62.
16. Mertz PM, Davis SC, Cazzaniga AL, Drosou A, Eaglstein WH. Barrier and antibacterial properties of 2-octyl cyanoacrylate-derived wound treatment films. *J Cutan Med Surg* 2003;7:1-6.
17. I.D. Fraser, A.C. Goede. Suture less circumcision. *BJU International* 2002;90: 467-468
18. Hayashi Y, Kojima Y, Mizuno K, Kohri K. Prepuce: phimosis, paraphimosis, and circumcision. *ScientificWorldJournal*. 2011;11:289-301.

19. Quinn J, Wells G, Sutcliffe T, *et al.* A randomized trial comparing octylcyanoacrylate tissue adhesive and sutures in the management of lacerations. *JAMA* 1997;277:1527-30.
20. Matin SF. Prospective randomized trial of skin adhesive versus sutures for closure of 217 laparoscopic port-site incisions. *J Am Coll Surg* 2003;196:845-53
21. James M Elmore, Edwin A Smith, Andrew J Kirsch. Suture less circumcision using 2-octyl cyanoacrylate (Dermabond): appraisal after 18-month experience. *Urology* 2007;4:803-6
22. Zempsky WT, Grem C, Nichols J, Parrotti D. Prospective comparison of cosmetic outcomes of simple facial lacerations closed with Steri-Strips or Dermabond. *Acad Emerg Med* 2001;8:438-9.
23. Singer AJ, Nable M, Cameau P, Singer DD, McClain SA. Evaluation of a new liquid occlusive dressing for excisional wounds. *Wound Repair Regen* 2003;11:181-7.
24. Toriumi DM, O'Grady K, Desai D, Bagal A. Use of octyl-2-cyanoacrylate for skin closure in facial plastic surgery. *Plast Reconstr Surg* 1998;102:2209-19.
25. Navid Jallali, Aryn Haji, Christopher J.E. Watson. A Prospective Randomized Trial Comparing 2-Octyl Cyanoacrylate to Conventional Suturing in Closure of Laparoscopic Cholecystectomy Incisions. *Journal of Laparoendoscopic & Advanced Surgical Techniques* 2004;14: 209-11.